EDUCATIONAL COLLECTIONS

OF

Australian Plants,

UNDER THE AUSPICES OF THE

VICTORIAN GOVERNMENT,

ISSUED BY

BARON FERD. VON MUELLER, C.M.G., M.D., PH.D., F.R.S.,

GOVERNMENT BOTANIST.

FIRST PART.

BY AUTHORITY:

JOHN FERRES, GOVERNMENT PRINTER, MELBOURNE.

MDCCCLXXIII.

Pimelea phylicoides ;

MEISSNER,

Lehmann Plantæ Preissianæ, vol. ii. pag. 271. (1847.)

ORDER THYMELEÆ.

New South Wales, Victoria, South Australia.

Port Phillip.

Contributor : G. Luehmann.

Stipa semibarbata ;

R. BROWN,

Prodromus Floræ Novæ Hollandiæ, pag. 174. (1810.)

ORDER GRAMINEÆ.

Southern tracts of Australia.

Port Phillip.—Contributor : G. Luehmann.

Prostanthera lasianthos;

LABILLARDIÈRE,

Novæ Hollandiæ Plantarum Specimen, vol. ii. pag. 18, tab. 157. (1806.)

Labiate-tree.

Order LABIATÆ.

South-east Australia.

Upper Yarra.

Contributor: G. Luehmann.



Eucalyptus viminalis;

LA BILLARDIÈRE,

Novæ Hollandiæ Plantarum Specimen, vol. ii. pag. 12, tab. 151. (1806.)

ORDER MYRTACEÆ.

South Australia, Victoria, Tasmania, and New South Wales.

Port Phillip.—Contributor : C. Groener.

Anguillaria australis;

FERD. VON MUELLER,

Fragmenta Phytographiæ Australiæ, vol. vii. pag. 74. (1870.)

VARIETY DIOICA.

Anguillaria dioica, R. Brown, Prodromus Floræ Novæ Hollandiæ, pag. 273. (1810.)

ORDER LILIACEÆ.

Widely dispersed through extratropical and eastern tropical Australia.

Port Phillip.—Contributor : C. Groener.



Leptorrhynchus tenuifolius;

FERD. VON MUELLER,

Fragmenta Phytographiæ Australiæ, i. 52. (1858.)

ORDER COMPOSITÆ.

Victoria.

Plains on the Yarra.

Contributor : C. Walter.

Rubus parvifolius;

LINNÉ,

Species Plantarum, editio secunda, pag. 707. (1762.)

Native Raspberry.

ORDER ROSACEÆ.

South and East Australia, East Asia.

Dandenong.

Contributor: C. Walter.

Hypoxis glabella;

R. BROWN,

Prodromus Floræ Novæ Hollandiæ, pag. 289. (1810.)

ORDER AMARYLLIDEÆ.

South Australia, Victoria, Tasmania.

Port Phillip.—Contributor : C. Groener.

Gentiana saxosa ;

G. FORSTER,

Svensk Kongl. Vetenskaps Academiens Handinger 1777, pag. 183, tab. 5.

ORDER GENTIANEÆ.

South-east Australia, Tasmania, New Zealand, South America

Properties : tonic.

Upper Yarra.

Contributor : C. Walter.

Myosotis suaveolens;

POIRET,

Encyclopédie Méthodique, Supplement, vol. iv. pag. 44. (1816.)

ORDER ASPERIFOLIÆ.

South-east Australia.

Upper Yarra.

Contributor : C. Walter.

Utricularia lateriflora;

R. BROWN,

Prodromus Floræ Novæ Hollandiæ, 431. (1810.)

ORDER LENTIBULARINÆ.

Tasmania, Victoria, New South Wales.

Heathbogs on Curdie's River.

Contributor : Baron Von Mueller.

Statice Taxanthema;

Rœmer et Schultes Systema Vegetabilium, vi. 793. (1820.)

ORDER PLUMBAGINEÆ.

From Victoria and Tasmania through East Australia to New Caledonia, China and Japan.

Saline meadows at Circular Head.

Contributor : Baron Von Mueller.



Myriophyllum amphibium;

LA BILLARDIÈRE,

Plantarum Novæ Hollandiæ Specimen, vol. ii. pag. 70, tab. 220. (1806.)

ORDER HALORAGEÆ.

Victoria and Tasmania.

Curdie's River.

Contributor : Baron Von Mueller.

Pultenæa Gunnii;

BENTHAM,

Annalen des Wiener Museums, vol. ii. pag. 82. (1840.)

ORDER LEGUMINOSÆ.

Victoria and Tasmania.

Dandenong.

Contributor: C. Walter.

Caulinia prostrata;

FERD. VON MUELLER,

Fragmenta Phytographiæ Australiæ, vol. vii. pag. 128. (1871.)

Kennedya prostrata, R. Brown in W. T. Aiton's Hortus Kewensis, second edition, vol. iv. 299. (1812.)

ORDER LEGUMINOSÆ.

Widely diffused through extratropical Australia.

Port Phillip.—Contributor : G. Luehmann.

Platylobium formosum ;

SMITH,

Specimen of the Botany of New Holland, pag. 17, tab. 6. (1793.)

ORDER LEGUMINOSÆ.

South-east Australia.

Dandenong.

Contributor : C. Walter.

Bossiæa cinerea;

R. BROWN,

Aiton, Hortus Kewensis, second edition, vol. iv. pag. 268. (1812.)

ORDER LEGUMINOSÆ.

South-east Australia.

Port Phillip.

Contributor : C. Groener.

Hovea heterophylla;

A. CUNNINGHAM,

J. Hooker, Flora Tasmanica, vol. i. pag. 93, tab. 15. (1860.)

ORDER LEGUMINOSÆ.

South-east Australia.

Grampians.

Contributor: D. Sullivan.



Leptospermum myrsinoides;

VON SCHLECHTENDAL,

Linnæa, xx. 653. (1847.)

ORDER MYRTACEÆ.

New South Wales, Victoria, South Australia.

Sandy heathland, Port Phillip.

Contributor : C. Grœner.

Anthistiria ciliata;

LINNÉ, FIL.,

Dissertatio Illustrans Nova Graminum Genera, p. 35. (1779.)

Kangaroo-Grass.

ORDER GRAMINEÆ.

Africa, South Asia, all Australia.

From Port Phillip.

Contributor : J. G. Luehmann.

Dicksonia antarctica;

LA BILLARDIÈRE,

Novæ Hollandiæ Plantarum Specimen, vol. ii. pag. 100, tab. 249. (1806.)

ORDER FILICES.

South-east Australia, New Zealand, Polynesia.

Upper Yarra.

Contributor: C. Walter.

Hakea nodosa;

R. BROWN,

In Transactions of the Linnean Society, x. 179. (1810.)

South Australia, Victoria.

From the Upper Yarra.

Contributor : C. Walter.



Pimelea humilis;

R. BROWN,

Prodromus Floræ Novæ Hollandiæ; 361. (1810.)

ORDER THYMELEÆ.

New South Wales, Victoria, South Australia, Tasmania.

From Port Phillip.

Contributor : C. A. Groener.

Convolvulus erubescens;

SIMS,

Botanical Magazine, t. 1067. (1808.)

ORDER CONVOLVULACEÆ.

Extratropic and Subtropic Australia.

From Port Phillip.

Contributor : J. G. Luehmann.

Helichrysum obcordatum;

FERD. VON MUELLER,

In Bentham's Flora Australiensis, iii., 632. (1866.)

ORDER COMPOSITÆ.

From the Loddon.

Contributor : Baron von Mueller.

Eucalyptus Stuartiana;

FERD. VON MUELLER,

In Nederl. Kruidk. Archiv., iv., 131. (1855.)

ORDER MYRTACEÆ.

New South Wales, Victoria, Tasmania, South Australia.

From Dandenong.

Contributor : Baron von Mueller

Melaleuca squarrosa ;

SMITH,

In the Transactions of the Linnean Society, vi. 300. (1802.)

ORDER MYRTACEÆ.

South-east Australia and Tasmania.

From the Yarra.

Contributor : C. Walter.



Drosera auriculata ;

BACKHOUSE,

Brogniart & Decaisne, Annales des Sciences Naturelles, Série troisième, vol. ix. pag. 295. (1848.)

Eared Sundew.

ORDER DROSERACEÆ.

From South Australia to the southern parts of Queensland, also in Tasmania and New Zealand.

Port Phillip.—Contributor : C. Groener.

Pelargonium Australe;

WILLDENOW,

Species Plantarum, iii., 675. (1800.)

ORDER GERANIACEÆ.

Extratropic Australia, New Zealand.

From Port Phillip.

Contributor: G. Luehmann.

Ranunculus hirtus;

BANKS AND SOLANDER,

In G. Forster, Florulæ Insularum Australium Prodromus, 90. (1786.)

ORDER RANUNCULACEÆ.

Victoria, New South Wales, New Zealand.

From the Yarra.

Contributor : C. Walter.

Acacia longifolia;

WILLDENOW,

Species Plantarum, vol. iv. pag. 1052. (1805.)

ORDER LEGUMINOSÆ.

From South Australia to South Queensland.

Gippsland.—Contributor : Baron von Mueller.

Drosera Whittackeri;

PLANCHON,

Annales des Sciences Naturelles, troisième série, vol. ix. pag. 302. (1848.)

ORDER DROSERACEÆ.

Victoria and South Australia.

Port Phillip.—Contributor: C. Richards.

Correa speciosa;

W. T. AITON,

Epitome of the Hortus Kewensis, pag. 366. (1814.)

ORDER RUTACEÆ.

From the Great Bight to South Queensland; also in Tasmania.

Port Phillip.—Contributor : C. Richards.

Pomaderris prunifolia ;

A. CUNNINGHAM,

Huegel, Enumeratio Plantarum Novæ Hollandiæ Austro-occidentalis, pag. 22. (1837.)

ORDER RHAMNACEÆ.

New South Wales, Victoria.

Upper Yarra.

Contributor : C. Walter.

Linum marginale;

ALL. CUNNINGHAM,

Hooker's London Journal of Botany, vol. vii. pag. 169. (1848.)

Native Flax.

ORDER LINEÆ.

Extratropic Australia.

Dandenong.

Contributor : C. Walter.

Comesperma ericinum;

CANDOLLE,

Prodromus Systematis Naturalis Regni Vegetabilis, vol. i. pag. 334. (1824.)

ORDER POLYGALEÆ.

South-east Australia.

Dandenong.

Contributor: C. Walter.

Billardiera cymosa;

FERD. VON MUELLER,

Transactions of the Victorian Institute 1855, 29.

ORDER PITTOSPOREÆ.

South Australia, Victoria.

You Yangs.

Contributor : J. P. Fullagar.

Stellaria flaccida ;

HOOKER,

Companion to the Botanic Magazine, vol. i. pag. 275. (1835.)

ORDER CARYOPHYLLEÆ.

Extratropic Australia.

Upper Yarra.

Contributor : C. Walter.

Hibbertia densiflora;

FERD. VON MUELLER,

Plants Indigenous to the Colony of Victoria, vol. i. pag. 15. (1860-62.)

ORDER DILLENIACEÆ.

South Australia, Victoria, New South Wales, Tasmania.

Port Phillip.

Contributor : G. Luehmann.

Aster orarius;

FERD. VON MUELLER,

Fragmenta Phytographiæ Australiæ, v. 78. (1865.)

Sea-coast Aster.

ORDER COMPOSITÆ.

Victoria, South Australia, Tasmania.

From Port Phillip.

Contributor: C. Groener.

Helipterum dimorpholepis ;

BENTHAM,

Flora Australiensis, iii., 650. (1866.)

ORDER COMPOSITÆ.

Extratropic Australia.

From Port Phillip.

Contributor : J. P. Fullagar.

Helichrysum obtusifolium ;

SONDER ET MUELLER,

Linnæa, xxv. 513. (1852.)

ORDER COMPOSITÆ.

Extratropical Australia.

Wilson's Promontory.

Contributor : Baron Von Mueller.

Tillæa intricata;

NEES VON ESENBECK,

Lehmann Plantæ Preissianæ, i. 278. (1845.)

ORDER CRASSULACEÆ.

Extratropical Australia, also in Tasmania.

Banks of the Loddon.

Contributor: Baron Von Mueller.

Centipeda Cunninghami ;

FERD. VON MUELLER,

Fragmenta Phytographiæ Australiæ, viii., 143. (1874.)

ORDER COMPOSITÆ.

Extratropic Australia.

From Dandenong.

Contributor : Baron von Mueller.

Clematis microphylla;

DE CANDOLLE,

Regni Vegetabilis Systema Naturale, vol. i. pag. 147. (1818.)

ORDER RANUNCULACEÆ.

Sparingly through the greater part of extratropical Australia.

Port Phillip.—Contributor : C. Schlipalius.